

2/5/04
CD

EA 1438


Document And Report Documentation Page Submitted as edoc_1075732948

Report Documentation Page		<i>Form Approved</i> OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.			
1. REPORT DATE 13 MAR 2003	2. REPORT TYPE N/A	3. DATES COVERED -	
4. TITLE AND SUBTITLE Detection and Classification from Hyperspectral Imagery Using the Normal Compositional Model		5a. CONTRACT NUMBER F19628-00-C-0002	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of the Defense		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited			
13. SUPPLEMENTARY NOTES Also see: ADM001520 , The original document contains color images.			
14. ABSTRACT			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:	17.	18.	19a. NAME OF RESPONSIBLE

a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	LIMITATION OF ABSTRACT UU	NUMBER OF PAGES 26	PERSON Patricia Mawby, EM 1438 PHONE:(703) 767-9038 EMAIL:pmawby@dtic.mil
----------------------------------	------------------------------------	-------------------------------------	---	--	---

Standard
Form 298
(Rev.
8-98)
Prescribed
by ANSI
Std
Z39-18

pwd: cannot determine current directory!



Detection and Classification from Hyperspectral Imagery Using the Normal Compositional Model

David Stein

ASAP 2003


11-13 March 2003

This work was sponsored by the Department of the Defense under Contract F19628-00-C-0002. Opinions, interpretations, conclusions, and recommendations are those of the author and are not necessarily endorsed by the United States government.

NCM_ASAP-1
DWJS 4/28/2003


MIT Lincoln Laboratory

20040317 140



Outline

- **Hyperspectral Imaging (HSI) aka Imaging Spectrometry**
- **Descriptive models of HSI**
- **The Normal Compositional Model**
- **Applications**
- **Summary**
- **Future Work**




Hyperspectral Imaging or Imaging Spectrometry


NASA: ER2 AVIRIS




NASA: E01 HYPERION




Hyperspectral Imager




NRL: HYDICE 0.4-2.4 nm



reflectance spectra of
various soils









Detection of a Known Target



RGB: HYPERION
(30m GSD,
195 bands down
sampled to 47,
400-2400 nm,
modeled with 10
background and
1 target classes)
subpixel
building

Detection statistic
(LMM target abundance)






HSI: Blind Unmixing

- Estimate class spectra from scene w/o library
- Estimate abundance of each material at each pixel

HYPERION RGB: 11 classes




Example class spectra




Abundance maps: water




Green vegetation







Outline

- **Hyperspectral Imaging (HSI) aka Imaging Spectrometry**
- **Descriptive models of HSI**
 - **Characteristics of HSI**
 - **Modeling intra-class variability**
 - **Common approaches to modeling HSI**
- **Applications**
- **Summary**
- **Conclusions**




Important Characteristics of HSI


**Different materials
occlude each other**




**Pixels are generally mixtures
(ARMY Night Vision Lab: NVIS-forest)**



AVIRIS: spectra from a roof





**Materials not well modeled by a
single spectrum**





Random Models of Spectral Variability: First and Second Order Statistics


**AVIRIS
(400-2400 nm)
Southern CA
multiple roof
reflectance
spectra**



**AVIRIS
(400-900 nm)
Tampa Bay
radiance spectra
identified with
phytoplankton
class**




- Correlation matrix is independent of radiance-to-reflectance transformation
- Correlation matrix is class dependent
- Significance of modeling variability judged by impact on performance




Subspace Models of Spectral Variability

- Subspace model:
 - Define a low-dimensional subspace such that target signatures may be replaced, with bounded error, by projection onto subspace
- Eigenvector construction
 - Given observations $\{x_1, \dots, x_m\} \subset R^n$ define $T = \sum_{i=1}^m x_i \cdot x_i^* = UDU^*$
 - Relative magnitude of the error vector obtained by ignoring the last N-p eigenvectors, where $N=\text{rank}(T)$, is

$$err^2(p) = \max_{j=p+1}^N \frac{|u_j \cdot x_i|^2}{|x_i|^2}$$
 - Applied to the roof data above




Dimension (p)	$err^2(p)$
5	0.0009
10	0.0001




Convex Models of Spectral Variability

- Find simplex such that every target is approximately represented as a convex mixture (linear combination such that coefficients are positive and sum to 1) of the vertices
- Construction of maximum volume inscribed n-simplex
 - Project samples onto first n eigenvectors of correlation matrix
 - Select n+1 affine independent samples
 - apply determinant update equations to maximize volume

**AVIRIS
roof data**



* Endmember
o Projected data



Common Approaches to Modeling HSI

- Local Normal model

$$x \in \text{Neigh}(z) \Rightarrow x \sim N(\mu_z, \Gamma_z)$$

- Normal mixture model

$$x \sim \sum_{j=1}^m \rho_j N(\mu_j, \Gamma_j), \rho_j \geq 0 \text{ and } \sum_{j=1}^m \rho_j = 1$$


- Subspace models (linear)

$$x_i = A\alpha_i + n, n \sim N(\mu, \Gamma)$$

- Linear mixture models (convex)

$$x_i = \sum_{j=1}^m a_{ij} s_j + n; a_{ij} \geq 0 \text{ and } \sum_{j=1}^m a_{ij} = 1; n \sim N(\mu, \Gamma)$$

- None of these models accounts for 1) occlusion, 2) intra-class variability, and 3) subpixel mixing



Outline

- Hyperspectral Imaging (HSI) aka Imaging Spectrometry
- Descriptive models of HSI
- The Normal Compositional Model
- Applications
 - Classification
 - Detection
- Summary
- Future Work



Normal Compositional Model

- Observation \vec{x}_i is modeled as

$$\vec{x}_i = \vec{e}_0 + \sum_{j=1}^m a_{ij} \vec{e}_j \text{ such that } \vec{e}_0 \sim N(\vec{\mu}_0, \Gamma_0) \text{ and } \vec{e}_j \sim N(\vec{\mu}_j, \Gamma_j)$$


subject to constraints

$$0 \leq a_{ij} \text{ for } j \leq r, \text{ and either } \sum_{j=1}^r a_{ij} = 1 \text{ or } \sum_{j=1}^r a_{ij} \leq 1; r \leq m.$$


- Features:


- Models subpixel mixing and random variation within a class
- Class parameters $\{(\mu_j, \Gamma_j) | 0 \leq j \leq m\}$ estimated as scene wide aggregates
- Estimates of class parameters converge under appropriate hypotheses to true values
- Abundance values $\{a_{ij} | 1 \leq j \leq m, 1 \leq i \leq N\}$ estimated at every pixel
- Additive components, e.g. noise and path radiance
- Accommodates (fat) subspace as well as (fat) simplex models.

- Special Cases: Linear mixture, Gaussian mixture and subspace models



NCM Blind Unmixing





NCM Detection

Inputs

Hyperspectral Image
 $\{x_i \mid 1 \leq i \leq N\}$

Number of Background Classes (m)

Target Models (t)
 $S', S'^{-1}, A', \{(\eta_\lambda, \Sigma_\lambda) \mid 1 \leq \lambda \leq t\}$

Outputs

NCM Background Parameter Estimator (scene average)
 $\{(\mu_j, \Gamma_j) \mid 0 \leq j \leq m\}$

Background Class Abundance Estimator (per pixel)
 $\{a_{ij} \mid 1 \leq i \leq N, 0 \leq j \leq m\}$

Background-plus-target Class Abundance Estimator
 $\{a_{ij} \mid 1 \leq i \leq N, 0 \leq j \leq m+t\}$

Decision Criteria

- 1) Anomaly
- 2) Likelihood Ratio
- 3) Abundance
- 4) Joint 2&3
- 5) Robust to target model uncertainties



Estimation of NCM Parameters: Nested Expectation Maximization

- Complete Likelihood function:

- N observations x_i and abundance vectors $\alpha_i = (a_{i1}, \dots, a_{im})$

$$p(x_1, \dots, x_N, a_{11}, \dots, a_{1m}, \dots, a_{N1}, \dots, a_{Nm} | \{(\mu_j, \Gamma_j) | 0 \leq j \leq m\}) = \prod_{i=1}^N N(x_i; \mu(\alpha_i) + \mu_0, \Gamma(\alpha_i) + \Gamma_0) p(\alpha_i)$$

- where

$$\mu(\alpha_i) = \sum_{j=1}^m a_{ij} \mu_j \text{ and } \Gamma(\alpha_i) = \sum_{j=1}^m a_{ij}^2 \Gamma_j$$

- Abundance values are hidden parameters

0. Initialize class parameters $\{(\mu_j^0, \Gamma_j^0) | 0 \leq j \leq m\}$

Linear mixture model techniques to identify initial endmembers (HSI)

Vertices of convex hull (low dimensional, e.g., multispectral, data)

1. Sample hidden parameters $\{a_{ij}^r | 0 \leq j \leq m, 1 \leq i \leq N\}$


Optimization of likelihood function (currently)

Monte Carlo Markov Chain (in progress)

2. Optimize class parameters for given values of hidden parameters

Expectation-Maximization Algorithm $\{(\mu_j^{i+1}, \Gamma_j^{i+1}) | 0 \leq j \leq m\}$

3. Repeat 1 and 2 until a convergence criterion is met



Updating Class Parameters Using Expectation Maximization

- Class parameters after iteration ℓ

$$\Omega^\ell = \{(\mu_j^\ell, \Gamma_j^\ell) \mid 0 \leq j \leq m\}$$


- Class mean update:

$$\begin{aligned}\mu_k^{\ell+1} &= \frac{1}{N} \sum_{i=1}^N E(e_k \mid x_i, \{a_{ij}^r\}, \Omega^\ell) \\ &= \mu_k^\ell + \frac{1}{N} \sum_{i=1}^N a_{ki}^\ell \Gamma_k^\ell [\Gamma_k^\ell(\alpha_i) + \Gamma_0^\ell]^{-1} (x_i - \mu^\ell(\alpha_i) - \mu_0^\ell)\end{aligned}$$

- Class covariance update


$$\Gamma_k^{\ell+1} = \frac{1}{N} \sum_{i=1}^N \text{cov}(e_k \mid x_i, \Omega^\ell) + [E(e_k \mid x_i, \Omega^\ell) - \mu_k^{\ell+1}] [E(e_k \mid x_i, \Omega^\ell) - \mu_k^{\ell+1}]^T$$

- Parameter updates are averages over expected values that are calculable from current parameters and abundance values



Outline

- Hyperspectral Imaging (HSI) aka Imaging Spectrometry
- Descriptive models of HSI
- The Normal Compositional Model
 - Generalization of common models
 - Estimation
 - Classification
 - Detection
- Applications
- Summary
- Future Work




AVIRIS Imagery of Cuprite, Nevada

AVIRIS: RGB Cuprite, NV




Acid sulfate hydrothermal alteration centers created 7.6-6.2 million years ago (hot sulfuric acid laden water flowing through surrounding rock changes the mineral content)

- Complex well studied scene used for evaluating algorithms
 - Mineral classification maps and spectral library available from US Geological Survey
 - USGS airborne hyperspectral identifications confirmed using ground spectrometry and laboratory analysis of field samples
 - 19 minerals plus 4 mixtures identifiable using SWIR data over 189 km² area
 - Subtle shifts in signatures due to variations in constituent elements, crystalline structure, temperature of formation
- Validate NCM estimation and blind unmixing
 - Sensor: AVIRIS
 - Spectral region: 50 bands (2-2.5 μ m)
 - Area covered: 42 km², 350 by 300 pixels extracted from a 189 km² image
 - Initial number of classes: 18



Identification of Spectra: Matching Absorption Features


• Alunite Group:




• Absorption Features:

- 2.17 μ : Al-O-H fundamental
- Higher formation temperature implies deeper and wider toward short end
- Shoulder: O-H stretch+Al-O-H bend
- Increasing concentration of Na shifts shoulder longer, and main band narrows
- 2.31 μ : O-H stretch+Al-O-H bend

Alunite Spectra from USGS Library



- NCM mean spectra identified with high temperature K-alunite (10) and moderate temperature mixed alunite (1) using USGS feature matching technique (correlation coefficients 0.99 and 0.98, respectively)






USGS Class Map and NCM Alunite Abundance Plane


NCM abundance: sum of classes 1&10



USGS Class Map




- Good qualitative comparison/ feature identification supports NCM unmixing
- Of 18 classes, 11 identified with minerals, 3 mixtures, 4 Fe bearing minerals





Detection Experiment: Life Vests in Ocean HSI

- Compare detection performance of Gaussian mixture, linear mixture, and NCM based known target and anomaly detection algorithms
- Background Data
 - 125-by-125 2 m² pixels
 - 24 band VNIR HSI (415-830 nm) from LASH sensor
- Target description
 - Life vest signature combined with background data from 1000 randomly selected pixels at 5% pixel fill fraction
 - 1 target class (mean given, covariance estimated as noise covariance)
- Model Parameters
 - 5 background classes




Scene RGB






Mean Class Spectra: Normal Mixture and Normal Compositional Models

Normal Mixture Model
(random, pure pixel)




No distinct sea-surface reflection class
formed using the normal mixture model


Normal compositional model
(random, mixed pixel)



NCM separates sea surface
reflections and upwelling light




Comparative Detection Performance





Summary and Conclusions

- Described a normal compositional model (NCM)
 - simultaneously treats mixed pixels and intra-class variability
 - accommodates subspace, convex, and random class variability models
 - generalizes and synthesizes normal mixture, linear mixture, and subspace models
- Applied NCM to Cuprite data
 - Class means identified with spectra of materials in scene
 - Abundance estimates qualitatively corresponded with USGS classification maps
- Application to ocean data
 - NCM estimation method identified classes where pure-pixel methods failed
 - NCM offered superior detection performance in comparison with LMM and GMM based models



Future Work

- Speed up the software
- Applications to real time HSI systems
- Applications to HSI performance models